



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Response to Letter: How Much Free Sugars Intake Should Be Recommended for Children Younger Than 2 Years Old?

Fidler Mis, Nataša ; Braegger, Christian ; Bronsky, Jiri ; Campoy, Cristina ; Domellöf, Magnus ;
Embleton, Nicholas D ; Hojsak, Iva ; Hulst, Jessie ; Indrio, Flavia ; Lapillonne, Alexandre ; Mihatsch,
Walter ; Molgaard, Christian ; Vora, Rakesh ; Fewtrell, Mary ; ESPGHAN Committee on Nutrition

DOI: <https://doi.org/10.1097/MPG.0000000000001803>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-165466>

Journal Article

Published Version

Originally published at:

Fidler Mis, Nataša; Braegger, Christian; Bronsky, Jiri; Campoy, Cristina; Domellöf, Magnus; Embleton, Nicholas D; Hojsak, Iva; Hulst, Jessie; Indrio, Flavia; Lapillonne, Alexandre; Mihatsch, Walter; Molgaard, Christian; Vora, Rakesh; Fewtrell, Mary; ESPGHAN Committee on Nutrition (2018). Response to Letter: How Much Free Sugars Intake Should Be Recommended for Children Younger Than 2 Years Old? *Journal of Pediatric Gastroenterology and Nutrition*, 66(3):e87-e88.

DOI: <https://doi.org/10.1097/MPG.0000000000001803>

REFERENCES

1. Zizzo AN, Valentino PL, Shah PS, et al. Second-line agents in pediatric patients with autoimmune hepatitis: a systematic review and meta-analysis. *J Pediatr Gastroenterol Nutr* 2017;65:6–15.
2. Sciveres M, Caprai S, Palla G, et al. Effectiveness and safety of ciclosporin as therapy for autoimmune diseases of the liver in children and adolescents. *Aliment Pharmacol Ther* 2004;19:209–17.
3. Czaja AJ. Autoimmune hepatitis: focusing on treatments other than steroids. *Can J Gastroenterol* 2012;26:615–20.
4. Alvarez F, Ciocca M, Cañero-Velasco C, et al. Short-term cyclosporine induces a remission of autoimmune hepatitis in children. *Hepatology* 1999;30:222–7.
5. Cuarterolo M, Ciocca M, Velasco CC, et al. Follow-up of children with autoimmune hepatitis treated with cyclosporine. *J Pediatr Gastroenterol Nutr* 2006;43:635–9.
6. Mennella JA, Bobowski NK, Reed DR. The development of sweet taste: From biology to hedonics. *Rev Endocr Metab Disord* 2016;17:171–8.
7. Welsh JA, Welsh JA1, Sharma A, et al. Consumption of added sugars and indicators of cardiovascular disease risk among US adolescents. *Circulation* 2011;123:249–57.
8. Harrison D, Stevens B, Bueno M, et al. Efficacy of sweet solutions for analgesia in infants between 1 and 12 months of age: a systematic review. *Arch Dis Child* 2010;95:406–13.
9. Vos MB, Kaar JL, Welsh JA, et al., American Heart Association Nutrition Committee of the Council on Lifestyle and Cardiometabolic Health; Council on Clinical Cardiology; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; Council on Functional Genomics and Translational Biology; and Council on Hypertension. Added sugars and cardiovascular disease risk in children. A scientific statement from the American Heart Association. *Circulation* 2017;135:e1017–34.

How Much Free Sugars Intake Should Be Recommended for Children Younger Than 2 Years Old?

To the Editor: A recent publication addressed the European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) Committee on Nutrition position about sugar in infants, children, and adolescents (1). Based on the World Health Organization guideline (2), the Committee stated that free sugar intake should be less than 5% of total calories for children and adolescents aged 2 to 18 years, suggesting that this figure should be lower to children younger than 2 years old. Establish a safe amount of sugar for infants may be a mistake because the evidence for this age group goes to the direction of nonexposure to the sweet taste early in life. Sweet taste experience in infancy triggers the preference for this flavor, which will determine children's poor dietary habits and consequently be a risk factor for noncommunicable diseases (3,4). Is the pediatric clinical use of sugar as an analgesic agent during painful procedures the main reason for the Committee to step away from a definite statement about sugar intake and infants (5)? Guidelines are intended for the general population and not for an individual one, which should be left open for the professional in clinical and technical responsibility. Therefore, based on evidence, the American Heart Association committee recommended (6): "Because there is minimal room for nutrient-free calories in the habitual diets of very young children, added sugars should be avoided in the diet of children <2 years of age". That is the statement we are expecting to be internationally widespread and hopefully help to build a healthy food environment for children.

Márcia Regina Vitolo

Graduate Program of Health Sciences, Universidade Federal de Ciências da Saúde de Porto Alegre Brazil

REFERENCES

1. Mis NF, Braegger C, Bronsky J, et al., ESPGHAN Committee on Nutrition. Sugar in infants, children and adolescents: a position paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 2017. [Epub ahead of print].
2. World Health Organization Guideline: Sugars Intake for Adults and Children Geneva, Switzerland: World Health Organization; 2015.

Response to Letter: How Much Free Sugars Intake Should Be Recommended for Children Younger Than 2 Years Old?

Reply: Dr Vitolo (1) asks what led the ESPGHAN CoN "to step away from a definite statement about sugar intake and infants" in our recent Position paper (2). We would like to clarify that our recommendation certainly does not mean that we are "stepping away" or establishing a "safe amount" of sugars for this age group.

The American Heart Association (AHA) recommends that "added" sugars should be avoided for children younger than 2 years (3). As explained in our article, the European Food Safety Authority (EFSA) and US definitions of "added sugars" (4,5) do not include sugars present in unsweetened fruit and vegetable juice and fruit juice concentrate, all of which are, however, captured in the WHO definition of free sugars (6), which we favour. Thus, children younger than 2 years who drink fruit juices and/or consume fruit juice concentrate, jellies, jams, preserves, and fruit spreads would still be following the AHA recommendation to avoid added sugars, whilst still consuming "free" sugars. Our recommendations state that "there is no nutritional requirement for free sugars in infants, children and adolescents" and suggest that the intake in those younger than 2 years should "probably be even lower" than the less than 5% of energy intake recommended for older children (2), partly considering the potential for sugar in infancy to influence taste and/or food preferences. We, however, did not provide a definitive figure because we found the evidence-base for doing so in this age group is not sufficiently robust, consisting largely of observational studies.

Furthermore, we have noted throughout the article the need to limit the intake of free sugars at all ages, and to clarify the terminology and labelling of sugars because consumers may not be aware that free sugars are present in foods and beverages, even when they are labelled as having "no added sugar" and "naturally occurring sugars." We hope that such measures would contribute to building a healthier food environment for infants and children.

ESPGHAN Committee on Nutrition

*Nataša Fidler Mis, †Christian Braegger, ‡Jiri Bronsky, §Cristina Campoy, ¶Magnus Domellöf, ¶Nicholas D. Embleton, #Iva Hojsak, **Jessie Hulst, ††Flavia Indrio,

^{††} ^{§§}Alexandre Lapillonne, ^{||||}Walter Mihatsch,
^{¶¶}Christian Molgaard, ^{##}Rakesh Vora, and ^{***}Mary Fewtrell
^{*}Department of Gastroenterology, Hepatology and Nutrition,
 University Children's Hospital, University
 Medical Centre, Ljubljana, Slovenia
[†]Division of Gastroenterology and Nutrition and Children's
 Research Center, University Children's Hospital,
 Zurich, Switzerland
[‡]Department of Paediatrics, University Hospital Motol, Prague,
 Czech Republic
[§]Department of Paediatrics, University of Granada,
 Granada, Spain
^{||}Department of Clinical Sciences, Pediatrics, Umeå University,
 Umeå, Sweden
[¶]Newcastle Neonatal Service, Royal Victoria Infirmary,
 Newcastle, UK
[#]University Children's Hospital Zagreb, Zagreb, Croatia
^{**}Erasmus MC, Sophia Children's Hospital, Rotterdam,
 The Netherlands
^{††}Ospedale Pediatrico Giovanni XXIII University of Bari,
 Bari, Italy
^{‡‡}Paris Descartes University, APHP Necker-Enfants Malades
 hospital, Paris, France
^{§§}CNRC, Baylor College of Medicine, Houston, TX
^{||||}Department of Pediatrics, Helios Hospital, Pforzheim, Germany
^{¶¶}Department of Nutrition, Exercise and Sports, University of
 Copenhagen, Hans Christian Andersen Children Hospital, Odense
 University Hospital, Odense, Denmark
^{##}Leeds Teaching Hospitals NHS Trust, Leeds, UK

^{***}Childhood Nutrition Research Centre, UCL GOS Institute of
 Child Health, London, UK

REFERENCES

1. Vitolo MR. How much free sugars intake should be recommended for children younger than 2 years old? *J Pediatr Gastroenterol Nutr* 2018; 66:e87.
2. Mis NF, Braegger C, Bronsky J, et al., ESPGHAN Committee on Nutrition. Sugar in infants, children and adolescents: a position paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 2017. [Epub ahead of print].
3. Vos MB, Kaar JL, Welsh JA, et al., American Heart Association Nutrition Committee of the Council on Lifestyle and Cardiometabolic Health; Council on Clinical Cardiology; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; Council on Functional Genomics and Translational Biology; and Council on Hypertension. Added Sugars and Cardiovascular Disease Risk in Children. A Scientific Statement From the American Heart Association. *Circulation* 2017;135:e1017–34.
4. EFSA: Review of labelling reference intake values. Scientific Opinion of the Panel on Dietetic Products, Nutrition and Allergies on a request from the Commission related to the review of labelling reference intake values for selected nutritional elements. *EFSA J* 2009;1008:1–14.
5. Institute of Medicine. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. Washington, DC: National Academies Press; 2005.
6. World Health Organization. Guideline: Sugars Intake for Adults and Children. Geneva, Switzerland: World Health Organization; 2015.